

~Since 1972~

## Corn & Soybean **Aerial Applied Insecticides**

INSECTICIDES						
Product	MOA	<b>Application Rates</b>	MIN GPA**	Compared To	PHI	REI
Carbaryl 4L	1A	0.5 to 1.5 qts./ac.	Aerial: 2; Ground: 10	Sevin® XLR	21 days	12 hou
Carbaryl 42.60%			tworm caterpillars, Dingy cutworm caterpillars, Green cloverworm caterpillars, Thistle caterpillars, Painted Yellowstriped armyworm caterpillars, Stinkbug nymphs and adults, and Bean leaf beetle adults.			
Pkg: 2.5 gals.	<b>Restrictions:</b> Do not apply more than 6 qts./ac. in a single season. Do not apply these products in combination with 2,4-DB he Avoid applications when bees are actively foraging.					
Chlorpyrifos 4E-AG*	1B	0.5 to 2 pts./ac.	Aerial: 2; Ground: 10	Lorsban® Advanced, Pilot™ 4E	28 days	24 hou
Chlorpyrifos 44.90%	caterpillars	, Yellowstriped armyworm catery	pillars, Grasshopper nymphs and	rootworm adults, Green cloverword adults, Soybean aphid nymphoeetle adults and bean leaf beetle a	s and adults	
Pkg: 2.5 gals., 265 gals.	nymphs and adults, Twospotted spider mite nymphs and adults, Japanese beetle adults and bean leaf beetle adults.  Restrictions: Do not make more than 3 applications in a single season. Do not apply more than 2 pts./ac. in a single application a do not reapply within 14 days. Do not apply more than 6 pts./ac. in a single season.					
Dimethoate LV-4	1B	19.1 fl. ozs./ac.; 0.5 lb. a.i./ac.; 1.0 to 1.5 pts./ac.	Aerial: 1; Ground: 10	Cygon®, Dimethoate 400	21 to 28 days	48 hou
Dimethoate 41.70%	Target Pes	1.0 to 1.5 pts./ac.		Cygon®, Dimethoate 400	28 days	48 hou adults, a
	Target Pes Bean leaf b	1.0 to 1.5 pts./ac.  tts: Grasshopper nymphs and adeetle adults.	lults, Soybean aphid nymphs and	75	28 days ymphs and	adults, a

Restrictions: Do not apply more than 0.03 lb. a.i./ac. in a single application. Do not reapply within 5 days. Do not apply more than

Target Pests: Black cutworm caterpillars, Dingy cutworm caterpillars, Green cloverworm caterpillars, Thistle caterpillars, Yellow striped

armyworm caterpillars, Grasshopper nymphs and adults, Soybean aphid nymphs and adults, Stinkbug nymphs and adults, Twospotted

Restrictions: Do not make more than 3 applications of Lambdafos or another product containing Chlorpyrifos in a single season. Do

not apply more than 38.0 fl. ozs./ac. in a single application. Do not reapply within 14 days. Do not apply more than 59.0 fl. ozs./ac. in

Target Pests: Fall armyworm caterpillars, True armyworm caterpillars, Yellowstriped armyworm caterpillars and Corn earworm

Restrictions: Do not apply more than 1 lb. a.i./ac. during a single application. Do not apply more than 2 lbs. a.i./ac. in a single season.

Cobalt® Advanced

Fyfanon®, Malathion 5

30 days

7 days

12 hours

Aerial: 2; Ground: 10

Aerial: 2; Ground: 10

spider mite nymphs and adults, Bean leaf beetle adults, Corn rootworm adults and Japanese beetle adults.

## Visit us at www.DrexChem.com for Specimens and SDS

0.06 lb. a.i./ac. in a single season.

1B; 3A 6 to 38 fl. ozs./ac.

16.0 to 25.6 fl. ozs./ac.;

Wait a minimum of 7 days between applications.

0.63 to 1 lb. a.i./ac.

a single season.

caterpillars.

Pkg: 1 gal., 2.5 gals.

Lambdafos™\*

Pkg: 2.5 gals., 265 gals.

**Malathion 5EC** 

Chlorpyrifos ...... 28.12%

Lambda-cyhalothrin ...... 1.44%

Malathion ..... 57.00%

(Corn only)

<sup>\*</sup>RESTRICTED USE PESTICIDE

<sup>\*\*</sup>GPA = Gallons per acre

The DREXEL logo and products marked with <sup>™</sup> and <sup>®</sup> are trademarks and registered trademarks of Drexel Chemical Company. All other brand names, product names or trademarks belong to their respective holders.

Always read and follow the label instructions on the product container(s) before using.

CSAAINSEC-0620

## **Drexel Sales Territories**

